



Steroidal saponins: An overview of medicinal uses

Zirwa Anwar¹ and Fatma Hussain^{1*}

¹Department of Biochemistry, Faculty of Sciences, University of Agriculture, Faisalabad-38040, Pakistan

Abstract

Modern study in drug discovery from medicinal plants is a versatile approach with a number of biological, phytochemical, molecular and botanical methods. Drug discovery is providing latest, significant and principal leads against various medical purposes like malaria, cancer, pain, HIV/AIDS and Alzheimer's disease. Numerous natural drugs of plant sources have either recently been purposed. Saponins are varied compounds broadly spread in the plant territory consist of steroid or triterpene aglycone and sugar chains. Saponins are commercially important compounds with increasing utilizations in cosmetics, pharmaceutical and food industries. Plant saponins have an extensive range of biological properties such as antifungal, antitumor, cytotoxic and anti-inflammatory.

Key words: Plants, saponins, antifungal, antitumor, cytotoxic

Full length article *Corresponding Author, e-mail: fatmauaf@yahoo.com

1. Introduction

Plants are consumed as antibiotic since ancient times. These antibiotics primarily acquire the appearance of unfinished drugs like that of tinctures, powders, poultices, teas and new herbal formulations. Extraction of active compounds that can be used as medicines from plants continues even today [1-6]. Minimum 7,000 medical compounds used in drug production in the modern array are derived from plants and 80% of them show optimistic relationship between new traditional use and therapeutic use of the plants from which they are extracted [7].

Saponin is a compound formed from a simple sugar. They consist of more than one sugar chains that may be present on a steroid and triterpenoidal aglycone frame. Backbone of saponin (aglycone) is known as sapogenin [8]. The functional groups depend upon the sapogenin, fraction of sugars; quality of the aglycone and nature of the sugars can differ significantly and conclude as very assorted compounds. The occurrence of saponins has been reported in more than 100 plants families [9, 10]. The saponin substance of plant resources is affected by the section of plant being observed, plant species, agronomic and environmental circumstances related to the plant growth and standard by-product analysis such as processing or storage. The term saponin is derivation of the Latin word "sapo" meaning soap [11, 12].

Steroidal saponins are normally occurring glycosides that have properties such as producing foam, complex formation with cholesterol. Steroidal saponins are found as plant secondary metabolites or a compound that helps proteins go through cellular membranes [13].

2. Chemical Composition

Steroidal saponins are extracted from the *Tribulus terrestris* plant. *Tribulus terrestris* contain a number of nutritional supplements along with 45% and 95% steroidal saponins. Protodioscin is the most important among them. Protodioscin is assumed to raise androgen receptor concentrations in cellular structures, encouraging those structures to become more susceptible to androgens like testosterone [14-18]. About 67 steroidal saponins were recognized on the basis of disintegration behaviors and reservation times of reference steroidal saponins (represented by sixty one peaks), including clusters of isomers. From them, fourteen saponins are new compounds [19]. Steroidal saponin are subgroup of steroids which are important class of organic compounds in which the substitution of cholesterol has modified to form 2 special frameworks, one is C-26 furostane (five ring structure) and another is C-27 spirostane (six ring structure, as dioscin). In case of spirostanols, spirokatal arrangement is linked at C-22 and sugar chain is attached at carbon-3. Furostanol has the skeleton such as that of spirostanol although in it sugar chain is not only linked to position C-3 but often also to carbon-26 and it is with open side chain [12].

3. Medicinal uses

Steroidal saponins could noticeably nourish the cardiac muscle, slow down platelet aggregation, increase coronary flow, improve peripheral circulation as well as show decline in cholesterol level and triglyceride in blood [13]. *Polygonatum* is a widely known popular Chinese medicine that is broadly used in Japan, Korea and China. In the last decade, it is reported that genus components have steroidal saponins, amino acids, lignins, alkaloids, flavones and carbohydrates, some of them show variable effects on the immune system [14]. Saponins like dioscin and its prosapogenins or gracillin have been recognized from yam. It has been accounted for diosgenin stifled cholesterol assimilation, expanded cholesterol discharge through biliary discharge, initiated segregation of lipoxygenase activity and furthermore actuated the programmed cell death process and cell cycle [15].

The rhizomes of *Dioscorea zingiberensis* are used for the isolation of diosgenin, sapogenin and its glycoside dioscin to produce steroid hormone, for example, sex hormone, progesterone and cortical hormone [20-22]. Then, steroidal saponins can likewise initiate troublesome responses, for example, normal weakness and cardiopalmus response [23]. The rhizome of *Anemarrhena asphodeloides* (Liliaceae) is used to prevent or reduce fever in traditional Chinese medicines. S. saponins are the better critical bioactive compounds in *A. asphodeloides*. It has antimicrobial action, represses carcinoma, diminishes radiation damage, minimizes the blood glucose level and is antagonistic to dementia [24-26]. Sarsapogenin and its proto structures are viewed as the important aglycones and ordinary sugars display in S. saponins are glucose, xylose and galactose [24, 27, 28]. Polyphyllin D, a steroidal saponin is present in *Paris polyphylla*, a customary Chinese therapeutic herb of Liliaceae family. *Paris polyphylla* displays spermicidal, hemostatic, antibacterial and pain relieving effects. The seed of *Paris polyphylla* are used for the treatment of inflammation, sour throat, abdominal pains, meningitis, respiratory tract tumors and leukemia [29-31].

A S. saponin isolated from *Agave attenuata* (Agavaceae) leaves inhibited increase in vascular permeability and destruction of red blood cells. Steroidal saponin from *Yucca schidigera* showed antifungal and antidermatophytic potency. A saponin division, containing principally mono desmosidic saponins had anti-yeast and antifungal action [32-34]. CAY-1, a S. saponin isolated from the products of *Capsicum frutescens* L. (Solanaceae) appeared to be an intense fungicide and anti-yeast agent [35]. A steroidal saponin, furcraestatin, in ethanolic leave extract of *Furcraea foetida* (L.) gave specific cytotoxicity in mouse fibroblasts [36]. S. saponins have both cytostatic and cytotoxic activities in leukemia [37-42]. Mimaki et al.

(1998) tested eleven new saponins from *Ruscus aculeatus* L. (Liliaceae). Only two, ruscogenin diglycoside (spirostanol saponin) and its relating 26-glycosyl-oxy-furostanol saponin indicated cell growth inhibitions. Mimaki et al. (1999) tested steroidal saponins of *Dracaena Draco* L. (Dracaenaceae) for their cytostatic activity. Only two saponins indicated cytostatic potential against the leukemia cells [36, 39].

Yucca schidigera is a Mexican herb with diverse therapeutic impacts. It has abundant steroidal saponins that may have antihypercholesterolemic and anti-arthritis potentials [43-46]. Various studies emphasized the influence of steroidal saponins on treatment of giardiasis, hypertension, hypercholesterolemia, arthritis, rheumatoid and malignant diseases [47-53]. S. saponins from *Paris polyphylla* have antitumor, antifungal and anthelmintic potentials [54-56]. Plant-determined items have indicated guarantee for the treatment across leishmaniasis [57]. Racemoside A, a water-solvent S. saponin refined from the products of shatavari, is a strong hostile to leishmanial particles [58].

4. Conclusions

Many plants contain rich source of S. saponins. Steroidal saponins isolated from different plants have many pharmacological uses. Therapeutic applications involved anti-inflammatory, antifungal, cytotoxic and antitumor, anti-arthritic, anthelmintic and anti leishmanial activities. Pharmaceutical industry now prefers the development of herbal medicines due to their easy availability and low cost.

References

- [1] G. Samuelsson. (2004). Drugs of Natural Origin: a Textbook of Pharmacognosy. 5th Swedish Pharmaceutical Press, Stockholm, Sweden.
- [2] M.J. Balick and P.A. Cox. (1997). Plants, People, and Culture: the Science of Ethnobotany. Scientific American Library, New York, USA.
- [3] A.D. Kinghorn. (2001). Pharmacognosy in the 21st century. Journal of Pharmacy and Pharmacology. 53(2): 135-148.
- [4] D.J. Newman., G.M. Cragg and K.M. Snader. (2000). The influence of natural products upon drug discovery. Natural Product Reports. 17(3): 215- 234.
- [5] M.S. Butler. (2004). The role of natural product chemistry in drug discovery. Journal of Natural Products. 67(12): 2141- 2153.
- [6] J.B. Marcy. (2005). Drug discovery from medicinal plants. Life Sciences. 78: 431- 441.
- [7] D.S. Fabricant and N.R. Farnsworth. (2001). The value of plants used in traditional medicine for drug discovery. Environmental Health Perspectives. 109: 69.

- [8] J. Bruneton. (1995). Pharmacognosy, Phytochemistry, Medicinal Plants. Lavoisier Publishing, Paris, France.
- [9] K.R. Price., I.T. Johnson and G.R. Fenwick. (1987). The chemistry and biological significance of saponins in foods and feeding stuffs. *CRC Critical Reviews in Food Science and Nutrition*. 26: 27-135.
- [10] K. Hostettmann and A. Marston. (1995). Saponins. Cambridge University Press. Cambridge, New York, USA.
- [11] G.R. Fenwick., K.R. Price., C. Tsukamoto and K. Okubo. (1991). Saponins. In: J.P.F. D'Mello, C.M. Duffus and J.H. Duffus, Eds. Toxic Substances in Crop Plants. The Royal Society of Chemistry, Cambridge, UK.
- [12] S.L. Deore., S. Khadabadi., S.K.P. Chittam., P.G. Bhujade., T.P. Wane., Y.R. Nagpurkar., P.D. Chanekar and R.G. Jain. (2009). Properties and pharmacological applications of saponins. *Pharmacology online Newsletter*. 2: 61-84.
- [13] N. Nakashima., I. Kimura., M. Kimura and H. Matsuura. (1993). Isolation of pseudoprotimosaponin AIII from rhizomes of *Anemarrhena asphodeloides* and its hypoglycemic activity in streptozotocin-induced diabetic mice. *Journal of Natural Products*. 56: 345-350.
- [14] D. Dinchev., B. Janda., L. Evstatieva., W. Oleszek., M.R. Aslani and I. Kostova. (2008). Distribution of steroidal saponins in *Tribulus terrestris* from different geographical regions. *Phytochemistry*. 69: 176-186.
- [15] R. Qureshi., G.R. Bhatti and R.A. Memon. (2010). Ethnomedicinal uses of herbs from northern part of Nara desert. Pakistan. *Pakistan journal of botany*. 42(2): 839-851.
- [16] H. Saima., B. Tamana., B.M. Khan and J. Asad. (2014). *Pakistan journal of botany*. 46(1): 399-404.
- [17] Z. Yang., K. Liping., Y. Heshui., Z. Jie., X. Chengqi., P. Xu., G. Yue., L. Chao and M. Baiping. (2013). Structure characterization and identification of steroidal saponins from the rhizomes of *Anemarrhena asphodeloides* by ultra-performance liquid chromatography and hybrid quadrupole time- of-flight mass spectrometry. *International Journal of Mass Spectrometry*. 7– 17: 341- 342.
- [18] Q. Sun., Y. Ju and Y. Zhao. (2002). Steroid saponins with biological activities. *Chinese Trad Herb Drugs*. 33: 276-280.
- [19] R. Yang., S. Tang., F. Pan., A. Zhao and Z. Pang. (2007). Advances in Study of *Dioscorea zingiberensis*. *Chinese Wild Plant Resources*. 26: 1-5.
- [20] State Pharmacopoeia Commission of the Ministry of Public Health. The pharmacopoeia of people's republic of China. China: 2000.
- [21] L. Nie., S. Lin and Z. Ning. (2004). Diosgenin in species of genus *dioscorea* research in progress. *Chinese Journal of Biochemical Pharmacology*. 25: 318-320.
- [22] L. Zan., W. Sun and K. Zhang. (2005). New technology study of saponin extract from *Dioscorea zingiberensis*. *Journal of Industrial and Engineering Chemistry*. 20: 138-139.
- [23] X.F. Zhang., Y. Cui., J.J. Huang., Y.Z. Zhang., Z. Nie and L.F. Wang. (2007). Immuno-stimulating properties of diosgenyl saponins isolated from *Paris polyphylla*. *Bioorganic & Medicinal Chemistry Letters*. 17: 2408-2413.
- [24] B. Lee., H.T. Trinh., K. Jung., S.J. Han and D.H. Kim. (2010). Inhibitory effects of steroidal timosaponins isolated from the rhizomes of *Anemarrhena asphodeloides* against passive cutaneous anaphylaxis reaction and pruritus, *Immunopharmacol Immunotoxicol*. 32: 357-363.
- [25] A. Tobari., M. Teshima., J. Koyanagi., M. Kawase., H. Miyamae., K. Yoza., A. Takasaki., Y. Nagamura and S. Saito. (2000). Spirostanols obtained by cyclization of pseudosaponin derivatives and comparison of anti-platelet agglutination activities of spirostanol glycosides. *European Journal of Medicinal Chemistry*. 35: 511-527.
- [26] Y.J. Kang., H.J. Chung., J.W. Nam., H.J. Park., E.K. Seo., Y.S. Kim., D. Lee and S.K. Lee. (2011). Cytotoxic, antineoplastic activity of timosaponin AIII for human cancer cells. *Journal of Natural Products*. 74: 701-706.
- [27] Y. Deng., B.P. Ma., Q.P. Xu., C.Q. Xiong., Z.Q. Liu., Y. Zhao and Y.C. Si. (2005). Effect and mechanism of effective component in Zhimu on ability of learning and memory in vascular dementia rats. *Chinese Pharmacological Bulletin*. 21: 833-837.
- [28] U.J. Youn., Y.S. Lee., H. Jeong., J. Lee., J.W. Nam., Y.J. Lee., E.S. Hwang., J.H. Lee., D. Lee., S.S. Kang and E.K. Seo. (2009). Identification of antiadipogenic constituents of the rhizomes of *Anemarrhena asphodeloides*. *Journal of Natural Products*. 72:1895-1898.
- [29] L.P. Kang., J. Zhang., Y. Cong., B. Li., C.Q. Xiong., Y. Zhao., D.W. Tan., H.S. Yu., Z.Y. Yu., Y.W. Cong., C. Liu and B.P. Ma. (2012). Steroidal glycosides from the rhizomes of *Anemarrhena asphodeloides* and their anti-platelet aggregation activities. *Planta Medica*. 78: 611-616.
- [30] B.P. da Silva., A.C. De Sousa., G.M. Silva., T.P. Mendes and J.P. Parente. (2002). A new bioactive

- steroidal saponin from *Agave attenuate*. *Zeitschrift für Naturforschung C*. 57: 423-428.
- [31] M. Miyakoshi., Y. Tamura., H. Masuda., K. Mizutani., O. Tanaka., T. Ikeda., K. Ohtani., R. Kasai and K. Yamasaki. (2000). Antiyeast steroidal saponins from *Yucca schidigera* (Mohave yucca). New anti-food-deteriorating agent. *Journal of Natural Products*. 63: 332-338.
- [32] V. Mshvildadze., A. Favel., F. Delmas., R. Elias., R. Faure., G. Decanosidze., E. Kemertelidze and G. Balansard. (2000). Antifungal and antiprotozoal activities of saponins from *Hedera colchica*. *Die Pharmazie*. 55: 325-326.
- [33] S.G. Sparg., M.E. Light and J. van Staden. (2004). Biological activities and distribution of plant saponins. *Journal of Ethnopharmacology*. 94: 219-243.
- [34] A. de Lucca., J.M. Bland., C.B. Vigo., M. Cushion., C.P. Selitrennikoff., J. Peter and T.J. Walsh. (2002). CAY-1, a fungicidal saponin from *Capsicum* sp. *Fruit. Medical Mycology*. 40: 131-137.
- [35] M. Itabashi., K. Segawa., Y. Ikeda., S. Kondo., H. Naganawa., T. Koyano and K. Umezawa. (1999). A new bioactive steroidal saponin, furcreastatin, from the plant *Furcraea foetida*. *Carbohydrate Research*. 323: 57-62.
- [36] Y. Mimaki., M. Kuroda., A. Kameyama., A. Yokosuka and Y. Sashida. (1998). Steroidal saponins from the underground parts of *Ruscus aculeatus* and their cytostatic activity on HL-60 cells. *Phytochemistry*. 48: 485-493.
- [37] Y. Mimaki., M. Kuroda., A. Kameyama., A. Yokosuka and Y. Sashida. (1998). Steroidal saponins from the rhizomes of *Hosta sieboldii* and their cytostatic activity on HL-60 cells. *Phytochemistry*. 48: 1361-1369.
- [38] Y. Mimaki., M. Kuroda., T. Asano and Y. Sashi. (1999). Triterpene saponins and lignans from the roots of *Pulsatilla chinensis* and their cytotoxic activity against HL-60 cells. *Journal of Natural Products*. 62: 1279-1283.
- [39] Y. Mimaki., M. Kuroda., A. Ide., A. Kameyama., A. Yokosuka and Y. Sashida. (1999). Steroidal saponins from the aerial parts of *Dracaena draco* and their cytostatic activity on HL-60 cells. *Phytochemistry*. 50: 805-813.
- [40] Y. Mimaki., A. Yokosuka., M. Kuroda and Y. Sashida. (2001). Cytotoxic activities and structure-cytotoxic relationships of steroidal saponins. *Biological and Pharmaceutical Bulletin*. 24: 1286-1289.
- [41] A. Yokosuka., Y. Mimaki and Y. Sashida. (2002). Spirostanol saponins from the rhizomes of *Tacca chantrieri* and their cytotoxic activity. *Phytochemistry*. 61: 73-78.
- [42] P.R. Cheeke., S. Piacente and W. Oleszek. (2006). Anti-inflammatory and anti-arthritis effects of *Yucca schidigera*: a review. *Journal of Inflammation*. 3.1: 6.
- [43] P.R. Cheeke. (1998). Natural toxicants in feeds, forages and poisonous plants. 2-Upper Saddle River. New Jersey: Prentice-Hall.
- [44] I.L. Lindhal., W.T. Shalkop., R.W. Dougherty., C.R. Thompson., G.R. Van Atta., E.M. Bickoff., E.D. Walter., A.G. Livingston., J. Guggolz., R.H. Wilson., M.B. Sideman and F. DeEds. (1957). Alfalfa saponins: Studies on their chemical, pharmacological and physiological properties in relation to ruminant bloat. USDA Technical Bulletin No. 1161. Washington. D.C
- [45] D. Oakenfull and G.S. Sidhu. (1989). Saponins. Toxicants of Plant Origin. Edited by: Cheeke PR. Boca Raton, Fla: CRC Press. 2: 97-141.
- [46] P. Griminger and H. Fisher. (1958). Dietary saponin and plasma cholesterol in the chicken. *Proceedings of the Society for Experimental Biology and Medicine*. 99: 424-426.
- [47] S.W. Kim., S.K. Park., S.I. Kang., H.C. Kang., H.J. Oh., C.Y. Bae and D.H. Bae. (2003). Hypocholesterolemic property of *Yucca schidigera* and *Quillaja saponaria* extracts in human body. *Archives of Pharmacal Research*. 26: 1042-1046.
- [48] I.T. Johnson., J.M. Gee., K.R. Price., C. Curl and G.R. Fenwick. (1986). Influence of saponins on gut permeability and active nutrient transport in vitro. *Journal of Nutrition*. 116: 2270-2277.
- [49] T.A. McAllister., C.B. Annett., C.L. Cockwill., M.E. Olson., Y. Yang and P.R. Cheeke. (2001). Studies on the use of *Yucca schidigera* to control giardiasis. *Veterinary Parasitology*. 97: 85-99.
- [50] N. Kaneda., H. Nakanishi and J. Staba. (1987). Steroidal constituents of *Yucca schidigera* plants and tissue cultures. *Phytochemistry*. 26: 1425-1429.
- [51] W. Oleszek., M. Sitek., A. Stochmal., S. Piacente., C. Pizza and P. Cheeke. (2001). Steroidal saponins of *Yucca schidigera* Roehl. *Journal of Agricultural and Food Chemistry*. 49: 4392-4396.
- [52] R. Bingham., D.H. Harris and T. Laga. (1978). *Yucca* plant saponin in the treatment of hypertension and hypercholesterolemia. *The Journal of Applied Nutrition*. 30: 127-136.
- [53] R. Bingham. (1976). New and effective approaches to the prevention and treatment of arthritis. *The Journal of Applied Nutrition*. 28: 38-47.
- [54] S.P. Zhang. (2007). Research progression chemical constituents and pharmacological effect of Genus *Paris*. *Strait Pharmaceutical Journal*. 19(16): 4-7.

- [55] J. Sun., B.R. Liu., W.J. Hu., L.X. Yu and X.P. Qian. (2007). In vitro anticancer activity of aqueous extracts and ethanol extracts of fifteen traditional Chinese medicines on human digestive tumor cell lines. *Phytotherapy Research*. 21:1102-1104.
- [56] Y.L. Fu., Z.Y. Yu., X.M. Tang., Y. Zhao., X.L. Yuan., S. Wang., B.P. Ma and Y.W. Cong. (2008). Pennogenin glycosides with a spirostanol structure are strong platelet agonists: structural requirement for activity and mode of platelet agonist synergism. *Journal of Thrombosis and Haemostasis*. 6: 524-533.
- [57] J. Dupouy-Camet. (2004). New drugs for the treatment of human parasitic protozoa. *Parasitologia*. 46: 81-84.
- [58] D. Avijit., G. Angana., M. Debayan., B.M. Nirup., B. Sukdeb., P.S. Niranjan and M. Chitra. (2007). *Journal of Medical Microbiology*. 56: 1196-1204.